



# Alcohol and Pregnancy

**If you drink alcohol during pregnancy, your baby may be at risk of lifelong birth defects.**

## Moderate Drinking: What's the Risk?

There is no safe amount or type of alcohol use during pregnancy. Even moderate drinking (one drink a day) can cause lifelong problems for your baby. These problems may be less obvious than those caused by heavy drinking. They may include problems with

- coordination
- behavior
- attention
- learning
- understanding consequences

## Heavy Drinking: What's the Risk?

Heavy drinking is having more than three drinks per occasion or more than seven drinks per week. The most severe result of heavy drinking during pregnancy is called fetal alcohol syndrome (FAS). FAS can cause serious birth defects for your baby, including

- problems with brain development
- lower-than-average height and weight
- smaller-than-normal head size
- abnormal facial features



## DID YOU KNOW?

- **No drinks are safe. One beer, one shot of liquor, one mixed drink, and one glass of wine all contain about the same amount of alcohol.**
- **If you are trying to get pregnant, you should not drink alcohol.**
- **Didn't know you were pregnant? While no amount or type of alcohol is safe during pregnancy, serious harm is unlikely if you drank before you knew you were pregnant. The most important thing is to stop drinking alcohol when you find out you are pregnant.**



**Alcohol-related birth defects are completely preventable.  
Do not drink alcohol during pregnancy.**

If it is hard for you to stop drinking, talk with your obstetrician–gynecologist (ob-gyn) or other health care professional about getting help. You also can visit the Alcoholics Anonymous website at [www.aa.org](http://www.aa.org) or call the Substance Abuse and Mental Health Services Administration's treatment referral line at 800-662-HELP (4357).

During your first prenatal visit, or at any time throughout your pregnancy, your ob-gyn or other health care professional can offer advice about avoiding alcohol while pregnant.

The American College of Obstetricians and Gynecologists believes that pregnant women who are dependent on alcohol should receive counseling and medical support to help them stop drinking.



The American College of  
Obstetricians and Gynecologists  
WOMEN'S HEALTH CARE PHYSICIANS  
409 12th Street SW, PO Box 96920  
Washington, DC 20090-6920  
[www.acog.org](http://www.acog.org)

PFS1015: This information was designed as an educational aid to patients and sets forth current information and opinions related to women's health. It is not intended as a statement of the standard of care, nor does it comprise all proper treatments or methods of care. It is not a substitute for a treating clinician's independent professional judgment. Please check for updates at [www.acog.org](http://www.acog.org) to ensure accuracy. Copyright December 2018 by the American College of Obstetricians and Gynecologists. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, posted on the internet, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission from the publisher.